

# LA-UR-22-20928

Approved for public release; distribution is unlimited.

**Title:** Local Time-Stepping: Matching Temporal Resolution to Spatial Resolution Highlight for LANL Institutional Computing

**Author(s):** Petersen, Mark Roger  
Capodaglio, Giacomo

**Intended for:** Report

**Issued:** 2022-02-03



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# Local Time-Stepping: Matching Temporal Resolution to Spatial Resolution

## Objective

- Develop an efficient parallel implementation and assess the performance of local time-stepping (LTS) schemes for the shallow water equations in the Model for Prediction Across Scales (MPAS).


## Approach

- Reformulate the LTS schemes to improve their computational performance while minimizing storage requirements.
- Use comparison with other established time-stepping schemes to validate the correctness of the implementation.
- Document the performance and computational timings of the LTS methods with respect to the state-of-the-art time integrator for the MPAS shallow water core.


## Impact

- It allows for faster simulations on existing variable-resolution meshes characterized by relatively few very high-resolution cells.
- The LTS capability improves the spatial resolution of existing variable-resolution meshes in localized areas.

Capodaglio G. and M. Petersen, "Local time stepping for the shallow water equations in MPAS," *J. Comput. Phys.*, **449**, 110818, (2022). [DOI: [10.1016/j.jcp.2021.110818](https://doi.org/10.1016/j.jcp.2021.110818)].



(Top) A diagram representing the different time-steps used on different regions of the computational grid. Blue region: fine time-step. All other regions: coarse time-step. (Bottom) The scalability of the LTS scheme is close to linear, only slightly worsening as the number of cores becomes large.



The order three LTS scheme can produce a CPU time savings of 70 percent for certain variable res meshes.